

AP Calculus

Unit 5 – Applications of the Derivative – Part 2

Day 7 Notes: Free Response Practice with Properties of Motion


NO CALCULATOR

A particle moves along the x – axis with velocity at time $t \geq 0$ given by $v(t) = -1 - e^{1-t}$.

- Find the acceleration of the particle at $t = 3$.
 - Is the speed of the particle increasing at $t = 3$? Give a reason for your answer.
 - Find all values of t at which the particle changes direction. Justify your answer.
 - The function $p(t) = e^{1-t} - t$ models the position of the particle for $t \geq 0$. Find the total distance that particle traveled on the time interval $0 \leq t \leq 3$.
-

NO CALCULATOR

A car is traveling on a straight road. For $0 \leq t \leq 24$ seconds, the car's velocity, $v(t)$, in meters per second, is modeled by the piecewise-linear function defined by the graph below.



- For what interval(s) of time does the car have zero acceleration? Show the work and explain the analysis that leads to your answer.
 - For each value of $v'(4)$ and $v'(20)$, find the value or explain why it does not exist. Indicate units of measure.
 - Let $a(t)$ be the car's acceleration at time t in meters per second per second. For $0 < t < 24$, write a piecewise-defined function for $a(t)$.
 - Find the average rate of change of v over the interval $8 \leq t \leq 20$. Does the Mean Value Theorem guarantee a value of c , for $8 < c < 20$, such that $v'(c)$ is equal to this average rate of change? Why or why not?
-

NO CALCULATOR PERMITTED

A particle moves along the x – axis so that its position at any time $t \geq 0$ is given by the function

$p(t) = t^3 - 4t^2 - 3t + 1$, where p is measured in feet and t is measured in seconds.

- a. Find the average velocity on the interval $t = 1$ and $t = 2$ seconds. Give your answer using correct units.
 - b. On what interval(s) of time is the particle moving to the left? Justify your answer.
 - c. Using appropriate units, find the value of $p'(3)$ and $p''(3)$. Based on these values, describe the motion of the particle at $t = 3$ seconds. Give a reason for your answer.
 - d. What is the maximum velocity on the interval from $t = 1$ to $t = 3$ seconds. Show the analysis that leads to your conclusion.
 - e. Find the total distance that the particle moves on the interval $[1, 5]$. Show and explain your analysis.
-

CALCULATOR PERMITTED

A test plane flies in a straight line with positive velocity $v(t)$, in miles per minute at time t minutes, where v is a differentiable function of t . Selected values of $v(t)$ for $0 \leq t \leq 40$ are shown in the table below

t (min)	0	5	10	15	20	25	30	35	40
$v(t)$ (miles per min)	7.0	9.2	9.5	7.0	4.5	2.4	2.4	4.3	7.3

- a. Find the average acceleration on the interval $5 \leq t \leq 20$. Express your answer using correct units of measure.
 - b. Based on the values in the table, on what interval(s) is the acceleration of the plane guaranteed to equal zero on the open interval $0 < t < 40$? Justify your answer.
 - c. Does the data represent velocity values of the plane moving away from its point of origin or returning to its point of origin? Give a reason for your answer.
 - d. The function f , defined by $f(t) = 6 + \cos\left(\frac{t}{10}\right) + 3\sin\left(\frac{7t}{40}\right)$, is used to model the velocity of the plane, in miles per minute, for $0 \leq t \leq 40$. According to this model, what is the acceleration of the plane at $t = 23$? What does this value indicate about the velocity at $t = 23$? Justify your answer, indicating units of measure.
-

AP Calculus AB
Unit 5 – Day 7 – Assignment

Name: _____

CALCULATOR PERMITTED

1) If $f(x) = \sin\left(\frac{x}{2}\right)$, then there exists a number c on the interval $\frac{\pi}{2} < x < \frac{3\pi}{2}$ that satisfies the conclusion of the Mean Value Theorem. Which of the following values could be c ?

- (A) $\frac{2\pi}{3}$ (B) $\frac{3\pi}{4}$ (C) $\frac{5\pi}{6}$ (D) π (E) $\frac{3\pi}{2}$

2) A particle moves along a line so that at time t , where $0 \leq t \leq \pi$, its position is given by $s(t) = -4\cos t - \frac{t^2}{2} + 10$. What is the velocity of the particle when its acceleration is zero?

- (A) -5.19 (B) 0.74 (C) 1.32 (D) 2.55 (E) 8.13